

1 1. A method for determining the concentration of chloride ions in samples,
2 comprising:

3 preparing an enzyme reagent, said enzyme reagent including:

4 α -amylase that is substantially calcium-free; and

5 an α -amylase activity detecting substrate; and

6 combining the enzyme reagent with sodium ion and a sample containing
7 chloride ion to be assayed, the sodium ion being present in a higher concentration
8 than said chloride ion;

9 assaying the quantity of α -amylase ^{activity} formed due to the presence of sodium ions
10 and chloride ions in said sample; and

11 determining the quantity of said chloride ions by reference to said assay of α -
12 amylase ^{activity}

13
14 2. The method according to claim 1, wherein calcium is removed from the α -
15 amylase that is substantially calcium-free by use of a chelating compound.

16
17 3. The method according to claim 1, wherein calcium is removed from the α -
18 amylase that is substantially calcium-free by use of a compound that forms a covalent bond
19 with calcium.

1 4. The method according to claim 2, wherein said chelating compound is a
2 member selected from the group consisting of ethylenediaminetetraacetic acid, trans-1,2-
3 cyclohexanediamine-N,N,N',N'-tetraacetic acid, glycol ether diamine tetraacetic acid,
4 iminotetraacetic acid, and diaminopropanetetraacetic acid.

5
6 5. The method of claim 2, wherein said chelating compound is
7 ethylenediaminetetraacetic acid.

8
9 6. The method according to claim 1, wherein said α -amylase activity detecting
10 substrate is a member selected from the group consisting of 2-chloro-4-nitrophenyl- α -D-
11 maltotrioxide, 4-nitrophenyl- α -D-maltopentaoside and α -glucosidase, 2-chloro-4-
12 nitrophenyl- β -D-maltopentaoside and α -glucosidase and β -glucosidase, 4-nitrophenyl- α -D-
13 maltoheptaoside, α -glucosidase, and 2-chloro-4-nitrophenyl- β -D-maltoheptaoside and α -
14 glucosidase and β -glucosidase.

15
16 7. The method according to claim 6, wherein said α -amylase activity detecting
17 substrate is 2-chloro-4-nitrophenyl- α -D-maltotrioxide.

18
19 8. The method according to claim 1, wherein said sample is a bodily fluid
20 sample.

21
22 9. The method according to claim 8, wherein said bodily fluid sample is
23 selected from the group consisting of serum, plasma, or urine.

24

1 10. The method of claim 1, wherein said sodium ion compound is sodium citrate.

2

3 11. The method of claim 1, wherein said sodium ion compound is sodium
4 acetate.

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

1 12. A composition for use in determining the concentration of chloride ions in a
2 fluid sample, comprising: α -amylase that is substantially calcium-free, sodium ion, and an α -
3 amylase activity detecting substrate.
4

5 13. A composition as in claim 12 further comprising a compound capable of
6 forming a chelate with a calcium ion and a calcium chelate compound.
7

8 14. A composition according to claim 13, wherein said compound capable of
9 forming a chelate with a calcium ion is a member selected from the group consisting of
10 ethylenediaminetetraacetic acid, trans-1,2-cyclohexanediamine-N,N,N',N'-tetraacetic acid,
11 glycol ether diamine tetraacetic acid, iminotetraacetic acid, and diaminopropanetetraacetic
12 acid.
13

14 15. A composition according to claim 13, wherein said compound capable of
15 forming a chelate with a calcium ion is ethylenediaminetetraacetic acid.
16

17 16. The composition according to claim 13, wherein said calcium chelate
18 compound is calcium-ethylenediaminetetraacetic acid.
19
20
21
22
23
24

1 17. The composition according to claim 12, wherein said α -amylase activity
2 detecting substrate is a member selected from the group consisting of 2-chloro-4-
3 nitrophenyl- α -D-maltotrioside, 4-nitrophenyl- α -D-maltopentaoside and α -glucosidase, 2-
4 chloro-4-nitrophenyl- β -D-maltopentaoside and α -glucosidase and β -glucosidase, 4-
5 nitrophenyl- α -D-maltoheptaoside, α -glucosidase, and 2-chloro-4-nitrophenyl- β -D-
6 maltoheptaoside and α -glucosidase and β -glucosidase.

7
8 18. The composition according to claim 12, wherein said α -amylase activity
9 detecting substrate is 2-chloro-4-nitrophenyl- α -D-maltotrioside.

10
11 19. The composition of claim 12, wherein said sodium ion compound is sodium
12 citrate.

13
14 20. The composition of claim 12, wherein said sodium ion compound is sodium
15 acetate.

16
17
18
19
20
21
22
23
24

1 (21.) A method of activating calcium-free α -amylase for enzymatic activity
2 comprising mixing chloride ion with calcium-free α -amylase in the presence of excess
3 sodium ion.

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

1 (22) A method for determining the concentration of sodium ions in samples,
2 comprising:

3 preparing an enzyme reagent, said enzyme reagent including:

4 α -amylase that is substantially calcium-free; and

5 an α -amylase activity detecting substrate; and

6 combining the enzyme reagent with excess chloride ion, and a sample
7 containing sodium ion to be assayed, the chloride ion being present in a higher
8 concentration than said sodium ion;

9 assaying the quantity of α -amylase formed due to the presence of sodium ions
10 and chloride ions in said sample; and

11 determining the quantity of said sodium ions by reference to said assay of α -
12 amylase.

13
14 23. The method of claim 22, wherein a calcium-binding compound is combined
15 with the enzyme reagent, the excess chloride ion, and the sample containing sodium ion to
16 be assayed before the α -amylase quantity is determined.

17
18 24. The method of claim 22, wherein said calcium-binding compound is
19 ethylenediaminetetraacetic acid.

20

21

22

23

24